

Orbital Functional Geometry XI

Schrödinger Equation in Orbital Space:
Linearisation, Orbital Potential, and von Mises Spectrum

Preprint

Judicael Brindel

Independent Researcher, Cotonou, Benin

ORCID: 0009-0007-4590-9874

March 2026

Abstract

We apply the Schrödinger equation to the Orbital Space $\mathcal{O}$ of *Orbital Functional Geometry I* (2026). The orbital Hamiltonian is $\hat{H}_{\mathcal{O}} = -(\hbar^2/2m)\Delta_{\mathcal{O}}$, where $\Delta_{\mathcal{O}}$ is the orbital Laplacian acting on $H_{\mathcal{O}} = \{f > 0, \text{ analytic on } S^1\}$ by $\Delta_{\mathcal{O}}f = (\ln f)''$.

The main result is an exact linearisation theorem: the substitution $u = \ln \psi$ transforms the orbital Schrödinger equation into the free Schrödinger equation on u :

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \Delta_{\mathcal{O}} \psi \quad \xrightarrow{u = \ln \psi} \quad i\hbar \frac{\partial u}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 u}{\partial \theta^2}$$

The nonlinearity disappears entirely in the natural coordinates of $\mathcal{O}$.

The stationary states are the von Mises functions $\psi_n^{(R,\phi)}(\theta) = e^{R \cos(n\theta - \phi)}$ with energy levels $E_n = \hbar^2 n^2 / 2m$ and infinite degeneracy parametrised by $(R, \phi) \in \mathbb{R}^+ \times [0, 2\pi)$.

The orbital potential — the effective potential seen by ψ in standard L^2 — is:

$$V_{\mathcal{O}}(\psi, \theta) = \frac{\hbar^2}{2m} \left(\frac{\partial_{\theta} \psi}{\psi} \right)^2 = \frac{\hbar^2}{2m} (\partial_{\theta} \ln \psi)^2$$

This is a state-dependent, positive-definite potential with no analogue in linear quantum mechanics.

This construction is distinct from the logarithmic Schrödinger equation (LogSE) of Rosen (1969) and Białynicki-Birula–Mycielski (1976): the LogSE remains non-linear in all coordinates, while the orbital Schrödinger equation linearises exactly under $u = \ln \psi$.

Contents

1	The Orbital Schrödinger Equation	4
2	Exact Linearisation	4
3	Stationary States and Spectrum	5
4	Temporal Dynamics	5
5	The Orbital Potential	6
6	Energy and Orbital Sobolev Norm	6

Introduction

The Orbital Space $\mathcal{O}$ constructed in Papers I–X carries a natural Hilbert structure $H_{\mathcal{O}}$, a self-adjoint Laplacian $\Delta_{\mathcal{O}}$, and a complete family of eigenfunctions (von Mises functions). These are the ingredients of a quantum mechanical system.

This paper applies the Schrödinger equation in $\mathcal{O}$ and derives its consequences. The main surprise is a complete linearisation: the orbital Schrödinger equation, nonlinear in ψ , becomes the standard free equation in $u = \ln \psi$.

This is not the logarithmic Schrödinger equation of Rosen (1969) and Białynicki-Birula–Mycielski (1976), which remains nonlinear in all coordinate systems and whose solutions are Gaussian (Gaussons). The orbital equation linearises exactly, and its solutions are von Mises functions — exponentials of cosines, not Gaussians.

1 The Orbital Schrödinger Equation

Definition 1 (Orbital Hamiltonian). *The orbital Hamiltonian is:*

$$\hat{H}_{\mathcal{O}} = -\frac{\hbar^2}{2m}\Delta_{\mathcal{O}}$$

where the orbital Laplacian acts on $\psi \in H_{\mathcal{O}}$ by:

$$\Delta_{\mathcal{O}}\psi = \frac{d^2}{d\theta^2} \ln \psi = (\ln \psi)''$$

Definition 2 (Orbital Schrödinger equation). *The orbital Schrödinger equation is:*

$$i\hbar \frac{\partial \psi}{\partial t} = \hat{H}_{\mathcal{O}}\psi = -\frac{\hbar^2}{2m}(\ln \psi)'' \psi$$

This is nonlinear in ψ , since $(\ln \psi)'' = \psi''/\psi - (\psi'/\psi)^2$.

2 Exact Linearisation

Theorem 1 (Linearisation theorem). *The substitution $u = \ln \psi$ transforms the orbital Schrödinger equation into the free Schrödinger equation:*

$$i\hbar \frac{\partial u}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 u}{\partial \theta^2}$$

The nonlinearity of the orbital equation disappears entirely in the natural coordinates of $\mathcal{O}$.

Proof. Set $u = \ln \psi$, so $\psi = e^u$ and $\partial_t \psi = \dot{u} e^u = \dot{u} \psi$. The orbital Schrödinger equation becomes:

$$i\hbar \dot{u} \psi = -\frac{\hbar^2}{2m} u'' \psi$$

Dividing by $\psi > 0$:

$$i\hbar \dot{u} = -\frac{\hbar^2}{2m} u'' \quad \square$$

Remark 1 (Why this differs from the LogSE). *The logarithmic Schrödinger equation (Rosen 1969; Białynicki-Birula–Mycielski 1976) has the form:*

$$i\hbar \partial_t \psi = -\frac{\hbar^2}{2m} \nabla^2 \psi + b(\ln |\psi|^2) \psi$$

This remains nonlinear under any substitution. Its solutions are Gaussian (Gaussons); it does not linearise. The orbital equation is a different object: it involves $(\ln \psi)''$ rather than $(\ln |\psi|^2)\psi$, and linearises exactly under $u = \ln \psi$.

Remark 2 (The variable $u = \ln \psi$ is the orbital invariant). *The variable $u = \ln \psi$ is precisely the orbital invariant $Q = \ln f/P - b$ of Paper I, restricted to $P = 1$, $b = 0$. The linearisation theorem is therefore not an accident: the natural coordinate of $\mathcal{O}$ is $\ln \psi$, and the dynamics are linear in that coordinate by construction.*

3 Stationary States and Spectrum

Theorem 2 (Stationary states). *The stationary states $\psi(\theta, t) = \psi_n(\theta)e^{-iE_n t/\hbar}$ satisfy $\hat{H}_O\psi_n = E_n\psi_n$, which in the u variable reads $u_n'' = -n^2u_n$.*

The solutions are $u_n(\theta) = R\cos(n\theta - \phi)$, giving:

$$\psi_n^{(R,\phi)}(\theta) = e^{R\cos(n\theta-\phi)}$$

These are the von Mises functions of Paper II.

Theorem 3 (Energy spectrum). *The energy levels are:*

$$E_n = \frac{\hbar^2 n^2}{2m}, \quad n = 0, 1, 2, \dots$$

with infinite degeneracy at each level $n \geq 1$, parametrised by $(R, \phi) \in \mathbb{R}^+ \times [0, 2\pi)$.

The ground state $n = 0$ has $E_0 = 0$ and $\psi_0 = \text{const}$ (unique up to normalisation).

n	E_n	Eigenfunction	Degeneracy
0	0	$\psi = c$	1
1	$\hbar^2/2m$	$e^{R\cos(\theta-\phi)}$	∞
2	$2\hbar^2/m$	$e^{R\cos(2\theta-\phi)}$	∞
n	$\hbar^2 n^2/2m$	$e^{R\cos(n\theta-\phi)}$	∞

Remark 3 (Comparison with the rigid rotor). *The rigid rotor has spectrum $E_n = \hbar^2 n^2/2I$ and eigenfunctions $e^{in\theta}$ (complex exponentials), with degeneracy 2 for $n \geq 1$ (states $\pm n$). The orbital spectrum has the same energy levels but different eigenfunctions (von Mises, not complex exponentials) and infinite degeneracy. The two systems have the same spectrum but different Hilbert structures.*

4 Temporal Dynamics

Theorem 4 (General time evolution). *The general state at $t = 0$ decomposes as:*

$$\psi(\theta, 0) = \exp\left(\sum_{n=1}^{\infty} R_n \cos(n\theta - \phi_n)\right)$$

by the decomposition theorem of Paper II. Its time evolution is:

$$\psi(\theta, t) = \exp\left(\sum_{n=1}^{\infty} R_n \cos\left(n\theta - \phi_n - \frac{\hbar n^2}{2m}t\right)\right)$$

Each mode n rotates its phase at angular velocity $\omega_n = \hbar n^2/2m$, quadratic in n .

Corollary 1 (Orbital dispersion). *Two modes n and m with equal initial phase $\phi_n = \phi_m$ dephase at rate $|\omega_n - \omega_m| = \hbar|n^2 - m^2|/2m$. Complete dephasing occurs at time:*

$$T_{nm} = \frac{2\pi m}{\hbar|n^2 - m^2|}$$

The dispersion is quadratic — faster than the linear dispersion of a free wave.

5 The Orbital Potential

Theorem 5 (Orbital potential). *In the standard $L^2(S^1)$ Hilbert space, the orbital Hamiltonian corresponds to the kinetic term plus an effective state-dependent potential. Comparing $\hat{H}_O\psi$ with $(-\hbar^2/2m)\partial_\theta^2\psi + V\psi$:*

$$V_O(\psi, \theta) = \frac{\hbar^2}{2m} \left(\frac{\partial_\theta \psi}{\psi} \right)^2 = \frac{\hbar^2}{2m} (\partial_\theta \ln \psi)^2$$

Proof.

$$-\frac{\hbar^2}{2m}(\ln \psi)''\psi = -\frac{\hbar^2}{2m} \left(\frac{\psi''}{\psi} - \left(\frac{\psi'}{\psi} \right)^2 \right) \psi = -\frac{\hbar^2}{2m}\psi'' + \frac{\hbar^2}{2m} \left(\frac{\psi'}{\psi} \right)^2 \psi$$

The first term is the free kinetic term; the second is $V_O\psi$. □

Remark 4 (Properties of V_O).

- **Positive definite:** $V_O \geq 0$ everywhere, with equality only where ψ is constant
- **State-dependent:** V_O depends on ψ itself — it is not a fixed external potential
- **Gradient energy:** V_O measures the squared logarithmic gradient of ψ — the cost of variation in orbital space
- **Invariant form:** $V_O = \frac{\hbar^2}{2m}(u')^2$ where $u = \ln \psi$ — it is the kinetic energy of u on S^1

Remark 5 (Relation to Bohmian mechanics). *The Bohmian quantum potential is $Q_{\text{Bohm}} = -(\hbar^2/2m)\nabla^2|\psi|/|\psi|$. The orbital potential $V_O = (\hbar^2/2m)(\nabla \ln \psi)^2$ has a similar structure — both are state-dependent and vanish for uniform ψ — but they are distinct objects. V_O is positive definite; Q_{Bohm} can take any sign. A precise comparison requires working with complex ψ , which is outside the scope of $\mathcal{O}$ as currently defined (where $\psi > 0$).*

6 Energy and Orbital Sobolev Norm

Theorem 6 (Energy as Sobolev norm). *The expectation value of energy in state $\psi = e^u$:*

$$\langle E \rangle = \frac{\hbar^2}{2m} \cdot \frac{1}{2\pi} \int_0^{2\pi} |u'(\theta)|^2 d\theta = \frac{\hbar^2}{2m} \|u\|_{H^1(S^1)}^2$$

The energy is the H^1 Sobolev norm of $u = \ln \psi$. It is minimised by $u = \text{const}$ (the ground state $\psi = \text{const}$).

Summary of New Results

Result	Statement	Status
Orbital Hamiltonian	$\hat{H}_{\mathcal{O}} = -(\hbar^2/2m)\Delta_{\mathcal{O}}$	Established
Linearisation	$u = \ln \psi$ gives free Schrödinger on u	Established
Distinction from LogSE	LogSE nonlinear; orbital equation linearises	Established
$u = \ln \psi$ is orbital Q	Linearisation follows from geometry of $\mathcal{O}$	Established
Stationary states	Von Mises functions $e^{R\cos(n\theta-\phi)}$	Established
Spectrum	$E_n = \hbar^2 n^2/2m$, infinite degeneracy	Established
vs. rigid rotor	Same spectrum, different eigenfunctions	Established
Time evolution	Phase rotation $\phi_n \rightarrow \phi_n + \omega_n t$, $\omega_n \sim n^2$	Established
Quadratic dispersion	Dephasing time $T_{nm} = 2\pi m/\hbar n^2 - m^2 $	Established
Orbital potential	$V_{\mathcal{O}} = (\hbar^2/2m)(\partial_{\theta} \ln \psi)^2$	Established
Positive definite	$V_{\mathcal{O}} \geq 0$, state-dependent	Established
Energy as H^1 norm	$\langle E \rangle = (\hbar^2/2m)\ u\ _{H^1}^2$	Established
Bohm relation	Similar structure, distinct objects	Established
Complex ψ extension	$\mathcal{O}$ currently requires $\psi > 0$	Open
Uncertainty principle	Non-canonical commutator, state-dependent bound	Open

Acknowledgements

This work is the direct continuation of the series *Orbital Functional Geometry* I–X (2026). The linearisation theorem was not anticipated: it emerged from applying the Schrödinger equation to $\mathcal{O}$ and following the substitution $u = \ln \psi$. The recognition that u is the orbital invariant Q of Paper I explained why the linearisation was inevitable. The distinction from the LogSE was verified by literature search before writing.

The author thanks Claude (Anthropic) for sustained mathematical dialogue throughout the development of this work.

References

- [1] J. Brindel, *Orbital Functional Geometry I–X*, Preprints, Cotonou, 2026.
- [2] G. Rosen, *Dilatation covariance and exact solutions in local relativistic field theories*, Phys. Rev. 183 (1969), 1186–1188.
- [3] I. Białynicki-Birula and J. Mycielski, *Nonlinear wave mechanics*, Ann. Phys. 100 (1976), 62–93.
- [4] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, Oxford University Press, 2nd ed., 1986.
- [5] D. Bohm, *A suggested interpretation of the quantum theory in terms of hidden variables*, Phys. Rev. 85 (1952), 166–179.